

Bion AA Forte

COMPOSITION

Component	Content
Free Amino Acids	7.8% w/w (10% w/v)
Total Nitrogen (N)	3.52% w/w (4.5% w/v)
Organic Nitrogen (N)	2.61% w/w (3.3% w/v)
Organic Matter	15.96% w/w (20.5% w/v)
Organic Carbon	8.78% w/w (11.3% w/v)
Presentation	Soluble Concentrate (SL)
Density	1.23 g/cc
pH (liquid solution)	5–6
pH (1% aqueous solution)	5–6
Color	Brown

CHARACTERISTICS

- Contains 100% active, free, and L-levorotatory amino acids
- Free amino acids that are codifiable and easily absorbed
- Stimulates vegetative growth and crop recovery
- Supports better development and productivity
- Promotes efficient absorption of fertilizers and micronutrients

DOSAGE AND INSTRUCTIONS FOR USE

Foliar Application:

Crop	Dose (cc/hl)	Instructions
Citrus, Olive, Vine, Fruit	150-300	2–3 applications during
Trees		flowering and fruit set
Vegetables	125-250	3–5 treatments every 10–15
		days after transplant or
		from 10-15 cm stage
Industrial Crops	150-250	2–3 applications during key
-		growth stages

APPLICATIONS TABLE

Crop	Dose (cc/hl)	PHI	
Apricot	150-300	N.A.	
Eggplant	125-250	N.A.	
Zucchini	125-250	N.A.	
Strawberry	125-250	N.A.	
Lemon Tree	150-300	N.A.	



Apple Tree	150-300	N.A.
Peach Tree	150-300	N.A.
Melon	125-250	N.A.
Orange Tree	150-300	N.A.
Olive Tree	150-300	N.A.
Other Citrus	150-300	N.A.
Pear Tree	150-300	N.A.
Pepper	125-250	N.A.
Grapefruit	150-300	N.A.
Watermelon	125-250	N.A.
Tomato	125-250	N.A.
Vine	150-300	N.A.

ADDITIONAL INFORMATION

Biostimulant Action:

Increases plant vigor and stimulates recovery.

Bioenergetic Effect:

Provides bioavailable amino acids essential for metabolism and protein synthesis.

Anti-Stress Protection:

Helps the plant resist adverse conditions: temperature, drought, salinity, pests, and diseases.

PRESENTATION

