

NatureFulvus 780

COMPOSITION

Component	Content
Organic Matter	99% w/v
Total Amino Acids	4.3% w/v
Fulvic Acids	78.10% w/v
Water-soluble Potassium Oxide (K ₂ O)	4.69% w/v
Organic Nitrogen (N)	2.0% w/v
Density	1.38 g/cc
рН	6–7
Electrical Conductivity (EC)	14.610 mS/cm
C/N Ratio	21.7

AMINOGRAM

Amino Acid	Percentage
Aspartic Acid	0.46%
Glutamic Acid	2.34%
Serine	0.19%
Proline	0.18%
Glycine	0.14%
Threonine	0.082%
Arginine	0.028%
Alanine	0.25%
Tyrosine	0.096%
Cystine	0.055%
Valine	0.15%
Methionine	0.027%
Isoleucine	0.25%
Phenylalanine	0.055%
Leucine	0.15%
Total Amino Acids	4.30%

CHARACTERISTICS

- Botanical extract-based product, concentrated, homogeneous, and naturally processed.
- Liquid organic concentrate that improves crop establishment and early plant development.
- Rich in fulvic acids with high root and vegetative growth promotion power.
- Fulvic acids complex micronutrients, improving their availability and absorption by crops.



APPLICATIONS TABLE

Crop	Dose	РНІ
Apricot	25-35 L/ha	N.A.
Eggplant	7–10 L/ha	N.A.
Zucchini	7–10 L/ha	N.A.
Plum	25-35 L/ha	N.A.
Lemon	25-35 L/ha	N.A.
Mandarin	25-35 L/ha	N.A.
Apple	25-35 L/ha	N.A.
Peach	25-35 L/ha	N.A.
Orange	25-35 L/ha	N.A.
Cucumber	7–10 L/ha	N.A.
Pear	25-35 L/ha	N.A.
Pepper	7–10 L/ha	N.A.
Grapefruit	25-35 L/ha	N.A.
Tomato	7–10 L/ha	N.A.
Vine	25-35 L/ha	N.A.

INSTRUCTIONS FOR USE

- Apply to the soil via fertigation, sprinkler, or flooding systems.
- Citrus, Fruit Trees, Vine: 25–35 L/ha split into 7.5 L/ha doses at vegetative growth, flowering, fruit set, and 20–30 days before harvest.
- Horticultural Crops: 7–10 L/ha every 15 days, especially in demanding phases, starting 15 days after transplant.
- Extensive Crops: 1.25–2.5 L/ha per application. Apply during early stages to promote root development.
- Use a minimum mix volume of 100 L/ha.
- Lawns: 7.5 L/ha after mowing. Apply in spring and autumn.
- Nurseries and Ornamentals: 7.5 L/ha every 7 days.
- Flood Irrigation: Apply at 30–40 L/ha per season.
- Foliar Application: Citrus, Olives, Vine, Fruit Trees (except plum): 150–300 cc/hl in 2–3 applications at pre-flowering, fruit set, and fruit development.
- Horticultural Crops: 125–250 cc/hl, 3–5 treatments every 10–15 days starting 1 week after transplant.
- Industrial Crops: 150–250 cc/hl, 2–3 treatments during the crop cycle.
- Apply NatureFulvus 780 foliar or via drip irrigation at critical crop stages or under stress: cold, heat, drought, salinity, pests, diseases, or nutritional deficiencies.

ADDITIONAL INFORMATION

- Botanical extract-based, concentrated, and homogeneous product obtained through natural processes.
- Improves crop establishment and early development.



- Rich in fulvic acids that enhance root and vegetative growth.
- Fulvic acids complex micronutrients, improving their availability and absorption by plants.

PRESENTATION

